

## COMPARISON OF MACROPHYTES DISTRIBUTION IN LAKE HAWASSA AT DIFFERENT POLLUTION ENTRY POINTS, HAWASSA, ETHIOPIA

LALISA GEMECHU BONA

College of Medicine and Health Science, Department of Environmental Health, Hawassa

University P.O. Box 1560 Hawassa, Ethiopia

Email: [laliga25@yahoo.com](mailto:laliga25@yahoo.com)

---

### Abstract

*Lake Hawassa is currently impinged by the various human activities in its watershed. This could create favourable conditions for the invasive macrophytes like water hyacinth attacking Lake Tana and Koka. Hence, identification of the impacts of pollution on the macrophytes species diversity, distribution and coverage is very important. In doing so, rapid bioassessment method in five pollution entry points, where major socio-economic activities prevail around the lake is applied. Thirty six samples were taken, following three transect lines, approximately equidistant from each other, and from each sampling site, onshore, up to the end of the macrophytes spreading into the lake, using rake. A total of 120 samples were collected and speciation was done using standard identification keys and photographs. Macrophytes coverage were approximated and different biotic indices were calculated for comparison between the sampling sites. The maximum distance of the macrophytes is 200m and minimum is 30m. The minimum distance was recorded around Tikur Wuha (30-100m); the only river flowing into the lake and maximum is at Amora Gedel (80-200m); the fish market area. The five species of macrophytes covering 8% of the lake are *Paspalidium germinatum*, *Nymphaea caerulea*, *Typha angustifolia*, *Potamogeton schweinfurthii*, and *Cyperus exaltatus*. The highest diversity ( $H'=1.25$ ) is observed around Dore Bafano site and lowest is at Tikur Wuha ( $H'=0.53$ ). The variation observed between these extreme Shannon diversity index values were ascribed to their relative species richness of Tikur Wuha and Dore Bafano ( $R=4$  and  $5$  respectively) and evenness ( $E=0.38$  and  $0.78$  respectively). There was greater organic pollution loads carried by the Tikur Wuha River when compared to other sample sites. On the other hand, all the sample sites have more than 75% Sorenson Similarity Index; indicating the impact of pollution between the sample sites is not that much different. Even though there are no invasive macrophytes species at this time, the disparity in diversity of the species supported at each specific site are one indication of the pollutants irregularly supporting the specific macrophytes. It is better to stop the pollutants prospectively than trying to abandon the invasive species after their existence.*

**Key Words:** *Pollutants, Macrophytes, Invasive Species, Community structure*

---

## Introduction

Freshwater Ecosystems have been critical to sustaining life and establishing civilizations throughout history (Small and Cohen, 1999). As indicated in Sudhira and Kumar (2000), Storm Water runoff and sewage discharges are the two common ways by which eutrophication and death of lakes happen (Sagneeta, 2007). Rapid population growth, urbanization, modern agriculture, poor infiltration when the adjacent land is concreted, commercial centers, Garages, land industrial development in the Hawassa City and its environs have posed potential risk to Lake Hawassa and its surroundings (Mulugeta, 2013).

According to Kibret and Harrison (1989), Lake Hawassa in comparison to the other Rift Valley Lakes, had a relative high water transparency and low sediment load because the large swampy sites to the east on the Tikur Wuha River called Cheleleka acts as efficient settling sites for silt coming from the higher hills and mountains of the eastern escarpment (Jacobus *et al.*, 2011). But this wetland is completely dry at currently (Forum for Environment, 2007). The surface of the Lake receives on average about 85 million m<sup>3</sup> of water annually through precipitation; however, the rate of open water evaporation on average, is about 160 million m<sup>3</sup>/year (Forum for Environment, 2007; Pattnaik, 2014). The loss of high amount of water through evaporation could be due to the extended coverage of macrophytes which may enhance the process to evapo-transpiration. There is a little data and information on trends of macrophytes abundance and distribution in Lake Hawassa (Pattnaik, 2014).

The Lake supports extremely poor women and children, who have no other income sources to support their families.

The water of the Lake is also used for bathing, recreation and as drinking water and other domestic uses, as well as for watering livestock and wildlife (Anasimos, 2013, Mulugeta, 2013). The fishery of the Lake supplies vital fish protein and incomes for the people of the sites and beyond. On the other hand, the trees of the relic forest around the Lake are inhabited by colorful birds chirping a lullaby with their mellifluous voice that titillate the hearts of people. Fishes dwelling in this Lake are food for the Plovers, Storks, Herons, Kingfishers and Fish Eagles (Mulugeta, 2013). Nowadays, the Lake Hawassa is seriously impinged by the various human activities in its watershed impacting water quality, storage capacity, recreational value, and natural lake bed habitat of natural and/or artificial lakes (Mulugeta, 2013).

## *The Role of Macrophytes in the Littoral Zone Ecosystem*

Macrophytes are the common features of an aquatic ecosystem. macrophytic vegetation plays an important role in maintaining the ecosystem of a lake (Sagneeta, 2007). Rooted macrophytes serve as a living link between the sediment, water, and (sometimes) atmosphere in wetlands, lakes, and rivers. The most notable function that plants serve is high primary production (Cronin *et al.*, 2006, Padial *et al.*, 2008). Besides, macrophytes are also involved in ecosystem processes such as bio-mineralization, transpiration, sedimentation, elemental cycling, materials transformations, and release of biogenic trace gases into the atmosphere (Cronin *et al.*, 2006).

The macrophytes vegetation serves as shelter, hiding and resting sites, and breeding ground for weed bed fauna (Annelids, Crustaceans and Insects), protozoans, rotifers, cladocerans,

copepods, ostracods and fish (Behailu *et al.*, 2010). Macrophytes and in particular *Potamogeton schweinfurthii* is the major food for the grazing hippopotamus in the lake as well as it is a shelter for these large mammals and for the different reptiles residing at the littoral region of the lake (Behailu *et al.*, 2010). On the other hand, it is important in curbing re-suspension of sediments and absorption of nutrients and toxic elements that have great impact on the lake water quality (Madsen, 2001).

The main threats to water quality in Africa include eutrophication, pollution, and the proliferation of invasive aquatic plants such as the water hyacinth (Ottichilo, 2003, Anderson, 2010) attacking Lake Tana and Koka of Ethiopia currently as it did in different parts of the world. Hence, quantifying the macrophytes coverage, distribution and diversity comparatively based on the adjacent socio-economic activities and identifying invasive species that could threaten the lake water quality and quantity is important to protect the lake.

#### **General Objectives**

- To identify the impact of pollutants on the macrophytes diversity, distribution and coverage in the Lake Hawassa and their respective pollution sources.

#### **Specific Objectives**

- To quantify the coverage of macrophytes in Lake Hawassa
- To identify the species of macrophytes in Lake Hawassa
- To compare biometric indices in terms of the selected adjacent socio-economic activities

### **Materials and Methods**

#### **Study Sites**

Lake Hawassa took its name from the City named Hawassa which is found at 270Km from the capital city of Addis

Ababa in Ethiopia. It is one of the rift valley lakes located at central location of Ethiopian rift valley at 275km south of Addis Ababa it is the smallest (96 km<sup>2</sup>) of the eight major lakes found in the rift valley of Ethiopia Mulugeta (2013) has found that the maximum depth of Lake Hawassa is 23.4m with an average depth of 11m. The lake drains an area of 1250 km<sup>2</sup>, and can accommodate  $1.036 \times 10^9$  m<sup>3</sup> of water. The major socio-economic activities adjacent to the lake are: the Fikir Hayk sites used mainly for recreation by food and drinking services; Amora Gedel sites, the major fish market place and storm water stabilization bond; Referral Hospital sites where hospital waste accumulation pond, storm stabilization pond is found; Dore Bafano where the adjacent land is used for crop production and Tikur Wuha sites where the only river carries different types of discharges to the lake.

#### **Study Design**

After receiving permission from the SNNPRS and NRMEPA of the City, Rapid assessment of macrophytes population in the five runoff entry points adjacent to the lake (Barbour *et al.*, 1999) following three equidistant transect lines perpendicular to the shore in the sites of interest (Fikir Hayk, Amora Gedel, Referral Hospital, and Dore Bafano) at three proportional spots per each transect. Each sampling spot provide composite of four pole-mounted rake (double rake head, 13.8 inch, with 14 teeth on each side) samples of macrophytes collected on the four directions perpendicular to. Hence, 12 rake samples were collected from each transect and 36 rake samples were collected from each sampling site. After pulling the rake from bottom-up direction, those macrophytes found in the sample were put in a plastic bag for identification

and quantification offshore using standard keys and check lists. macrophytes coverage was estimated by the interval between the longest and shortest transect lines at each sites and five key informants about the presence of gradual change on macrophytes coverage overtime were asked.

The recorded data were averaged in their species category for each site and tabulated for comparison. In addition, diversity and similarity indices were calculated specifically for each species in the sampling sites and comparatively between sites to identify the variability between species composition, distribution enabling their difference indicating the impact of pollutants in the sampling sites. During data collection, trained data collectors were provided with rake, meter, boat, hat, protective cloths, and life-saving jacket, glove and first aid kits, log books, pens, markers, plastic bags, Global Positioning System and digital camera.

**Results and Discussions**

**Macrophytes Coverage**

The macrophytes cover of the littoral zone stretched out up to 200 m (30–200m) (table 2). The overall macrophytes cover decreased in the order of: Amora Gedel (80–200m) > Fikir Hayk (80–140m) > Referral Hospital (50–160m) > Dore Bafano (80–90m) > Tiku Wuha (30–100m). Preceding studies (Tadesse, 2005; Behailu *et al.*, 2010) on the Lake Hawassa found that the littoral sites extends out to 150 m off shore and roughly follows the 4 m isobaths. Likewise, Amare (2005) that the littoral zone of Lake Hawassa is fringed by an emergent and submerging macrophytes belt that extends offshore down to 4m depth. By all appearances, the macrophytes cover observed in study modest increment; even though the key informants there is no change in the

coverage and it is the water which is increasing on the littoral zone.

Table 1: Macrophytes distribution in selected littoral sites of Lake Hawassa

Sampling sites	Transect lines (m)	
	Max.	Min.
Amora Gedel	80	200
Fikir Hyk	80	140
Tikur Wuha	30	100
Dore Bafano	80	90
Referral Hospital	50	160

Taking a rough assumption of the 200m off shore distance of the macrophyte coverage by assuming circular shape for the Lake (the lake have the longest distance of 16km and shortest 9km averaged to 12.5km diameter) the macrophyte coverage could be calculated as;  $\Pi(R^2-r^2)$ , Where R is the radius of the lake (12.5/2km) and r (200m=0.2km) is the distance of macrophytes from the shore. Hence macrophytes coverage is approximately 39.1km<sup>2</sup> which is roughly 8% of the lake surface. This is very few when compared to Lake Tana (89%). Added up on the disappearance of Cheleleka Wetland (Yezbie, 2016), which was acting as a lung of the lake, the water quality be maintained by the existing low coverage and few species. Even though macrophytes maintain the clear-water state by a variety of mechanisms (e.g., stabilizing sediments, promoting zooplankton populations) (Cronin *et al.*, 2006).

**Macrophyte species**

The species identified over the five sampling stations were *Paspalidium germinatum*, *Nymphaea caerulea*, *Typha angustifolia*, *Potamogeton schweinfurthii*, and *Cyperus exaltatus* (Table 2). All of the macrophytes were emergent or floating; where there are submerged. previous studies on lake Hawassa reported that the

littoral zone is covered with the various emergent, submerged and floating wetland flora including *Nymphaea caerulea*, *Potamogeton schweinfurthii*, *Typha angustifolia*, *Paspalidium germinatum* (Tadesse, 2005; Behailu *et al.*, 2010; Pattnaik, 2014), *Cyperus exaltatus* (Tadesse, 2005; Behailu *et al.*, 2010) and *Ludwigia stolonifera* (Behailu

*et al.*, 2010; Pattnaik, 2014). Hence, there no invasive species like water hyacinth in the macrophytes species of the lake not identified in this study. But the very few compared to Lake Tana; which comprise about 71 different species of macrophytes (Yesbie Kassa, 2016).

Table 2: Macrophytes composition, frequency of occurrence and abundance

Macrophytes species	Total		Fikir Hayk		Amora Dedel		Referral Hospital		Tikur Wuha		Dore Bafano	
	Freq.	Abundance	Freq.	Abundance	Freq.	Abundance	Freq.	Abundance	Freq.	Abundance	Freq.	Abundance
<i>Paspalidium germinatum</i>	100	296.4	100	93.3	100	75.7	100	61.0	100.0	142.7	100	121.3
<i>Nymphaea caerulea</i>	100	97.6	100	84.3	100	40.7	33.3	18.0	33.3	18.0	66.6	38.5
<i>Typha angustifolia</i>	80	117.5	33.3	101.0	66.6	43.0	66.6	24.5	0.0	0.0	100	78.0
<i>Potamogeton schweinfurthii</i>	80	71.3	33.3	9.0	0	0.0	100	40.7	66.6	25.0	100	34.7
<i>Cyperus exaltatus</i>	80	6.8	33.3	7.0	33.3	9.0	0	0.0	33.3	5.0	33.3	6.0

*Paspalidium germinatum* and *Nymphaea caerulea* were found in all the five stations while the other three macrophytes were recorded from only four sampling sites. On the other hand, *Paspalidium germinatum* observed at every sub-station. Behailu *et al.* (2010) observe that *Paspalidium germinatum* the dominant macrophytes of the lake. Apart from this, *Potamogeton schweinfurthii*, *Typha angustifolia* and *Cyperus exaltatus* absent in Amora Gedel, Tikur Wuha, and Referral Hospital stations. Hence, even though have large macrophytes coverage, it hold true for the evenly distribution of each species. The corresponding data on abundance presented in the selfsame table reveals that the most frequent *Paspalidium germinatum* also the most dominant macrophytes across the sampling stations. Besides, *Typha angustifolia* followed by *Potamogeton*

*schweinfurthii* considerable abundance over three and two stations, respectively. Apart from this, Behailu *et al.* (2010) that the inflow sub-zone at the mouth of Tikur Wuha river has typical macrophytes consisting of large patches of *Cyperus papyrus*, *Lemna minor*, *Wolffia arrhiza* and *Pistia stratiotes*.

#### **Shannon diversity index**

Shannon diversity index, which characterize biotic integrity and natural balance between flora and fauna without specifically diagnosing ecological conditions and causes of impairment (Barbour *et al.*, 1999) indicate across the sampling stations ranged from the lowest value ( $H' = 0.53$ ) at Tikur Wuha highest value ( $H' = 1.25$ ) at Dore Bafano (Table 3). But 89% of the species found in these sample sites from the calculated Sorenson's similarity index (Table 4) and the variation could be due to the few

number of taxa (n=5 from table 1) identified does not fulfil the minimum number of taxa recommended (n=10) by Barbour et.al, 1999. The variation observed between these extreme Shannon diversity index values ascribed to their relative between Tikur Wuha and Dore Bafano (R=4 and 5 respectively) and (E=0.38 and 0.78 respectively). Species richness is predicted to decrease with increasing pollution because many species are stressed (Barbour et.al, 1999) an indication of serious entry of pollutants on Tikur Wuha side of the lake; mainly by the Tikur Wuha river carrying industrial effluents and other anthropogenic wastes (mainly organic in their origin according to Barbour et.al, 1999) . There is a substantial difference of diversity, on the shore lines of each sampling spots transect lines than at the middle and at the edge of the

The analysis of the macrophytes species with recourse to Sorenson index of similarity that there marked similarities ( $\geq 0.75$  or  $\geq 75\%$ ) between the sampling stations (Table 4). Even though there is some diversity difference between Tikur Wuha and Dore Bafano, the difference arise only the Evenness not from species Richness as indicated previously. This is also justified by species Similarity between Dore Bafano and Fikir Hayk (SR=5) while the Evenness is different (Table 4 above). This, in turn to the absence of species like *Typha angustifolia*

in Tikur Wuha, *Potamogeton schweinfurthii* in Amora Gedel and *Cyperus exaltatus* Referral Hospital. This has determined the less similarity (75%) of these sites among themselves compared to the other sites. In line with the types and sources of pollutants, these sites are contributing a lot concentrated point source wastes from hospital and storm water stabilization ponds (Referral Hospital site), industrial wastes and wastes from different sources including the storm water entering into the lake carried by the only tributary called Tikur Wuha (at Tikur Wuha site) and wastes from the fish market, fish food processing houses, storm water stabilization pond (Amora Gedel site). While there is waste from the recreational sites of Fikir Hayk and the adjacent storm water accumulation artificial wet lands (Fikir Hayk site) and solely agricultural wastes from Dore Bafano Site.

Better macrophyte diversity indicates the relative health of the ecosystem at that specific site while specific species found in that site could infer some additional information for comparing the types of pollutants entering into the lake at each study sites. *Typha angustifolia* does not prefer chemical pollutants like Cadmium and Lead (Klaudia, 2016); the reason it could be absent from the Tikur Wuha site and indicating not only the organic pollutants but also inorganic pollutants of anthropogenic origin is serious at this site.

Table 3: Diversity and of the macrophytes

Shannon diversity index	Fikir Hayk				Amora Gedel				Referral Hospital				Tikur Wuha				Dore Bafano			
	T	S	M	En	T	S	M	En	T	S	M	En	T	S	M	En	T	S	M	En
<b>H'</b>	1.13	1.23	0.68	0.69	1.09	1.18	0.51	0.91	1.13	1.08	0.50	1.23	0.53	0.69	0.26	0.28	1.25	1.28	1.10	0.94
<b>SE</b>	0.70	0.76	0.98	1.00	0.79	0.85	0.79	0.83	0.81	0.98	0.72	0.89	0.38	0.63	0.37	0.40	0.78	0.80	0.79	0.85
<b>SR</b>	5	5	2	2	4	4	2	3	4	3	2	4	4	3	2	2	5	5	4	3

H' = Shannon diversity index, SE= Species, SR= Species richness, T=total, S=shore, M=middle, En=end *Sorenson index of similarity and  $\beta$ -diversity*

Table 4: Sorenson index of similarity (s) and -diversity of the Macrophytes

Sorenson diversity index	Sample sites comparison										
		Fikir Hayk-Amora Gedel	Fikir Hayk-Referral Hospital	Fikir Hayk-Tikur Wuha	Fikir Hayk-Dore Bafano	Amora Gedel-Referral Hospital	Amora Gedel-Tikur Wuha	Amora Gedel-Dore Bafano	Referral Hospital-Tikur Wuha	Referral Hospital-Dore Bafano	Tikur Wha-Dore Bafano
A	5	5	5	5	4	4	4	4	4	4	
B	4	4	4	5	4	4	5	4	5	5	
C	4	4	4	5	3	3	4	3	4	4	
Index of similarity (S) $S = 2C/A+B$	2C	8	8	8	10	6	6	8	6	8	8
	A+B	9	9	9	10	8	8	9	8	9	9
Similarity	$2C/A+B$	0.89	0.89	0.89	1.00	0.75	0.75	0.89	0.75	0.89	0.89

### Conclusions and Recommendations

The macrophytes distance of the littoral zone stretched out up to 200 m from the shore. The total coverage is about 39.1Km<sup>2</sup>; which is about 8% of the lake. It is maximum around Amora Gedel sites compared to the others; while least was at Tikur Wuha sites; which is further confirmed by the calculated diversity index at the same site and could be due to high inorganic and organic chemicals entering the lake from the Tikur Wuha site Tikur Wuha River. This can also be deduced from the characteristics of *Typha angustifolia*. The macrophytes were composed of five species namely *Nymphaea caerulea*, *Potamogeton schweinfurthii*, *Typha angustifolia*, *Paspalidium germinatum*; which were very few compared to other lakes in Ethiopia. Even so, only two species (*Paspalidium germinatum* and *Nymphaea caerulea*) were common to in all the sites while *Potamogeton schweinfurthii*, *Typha angustifolia* and *Cyperus exaltatus* were absent in Amora Gedel, Tikur Wuha, and Referral Hospital stations. *Paspalidium germinatum* was the most abundant macrophyte in all the sampling sites.

Hence, control of pollutants to these sites is very crucial; paying attention to Tikur Wuha sites and followed by Referral Hospital, Amora Gedel, Fikir Hayk and finally Dore Bafano.

### References

- Amare, L. (2005). Site action plan for the conservation and sustainable use of the Lake Hawassa biodiversity. Valley lakes project. Institute of Biodiversity Conservation.
- Anasimos, Tekalign (2013). *The Impact of Unplanned Urban Waterfront Development Lake Hawassa*. Ethiopian Institute of Architecture, Building Construction and City Development, Addis Ababa.
- Andersen, C. (ed). (2010). *Questioning Greater Yellowstone's Future: Climate, Land Use, and Invasive Species. Proceedings of the 10th Biennial Scientific Conference on the Greater Yellowstone Ecosystem*. Mammoth Hot Springs Hotel, Yellowstone National Park. WY, and Laramie.
- Barbour, M.T., Gerritsen, J., Snyder, B.D. and Stribling, J.B. (1999). Rapid Bioassessment Protocols for use in streams and Wadeable Rivers: Periphyton, Benthic Macroinvertebrates and Fish, second edition. U.S. Environmental Protection Agency; Office of Water. EPA 841-B-99-002. Washington, D.C.
- Behailu, B., Bekele, L., Yosef, T/G., Kefyalew, S., and Wondwossen, S. (2010). Environmental Impact Assessment and Policy on Lake Hawassa, SNNPRS. SOS-Sahel, Ethiopia
- Cronin, G., William, M., Lewis Jr., and Michael, A.S. (2006). Influence of freshwater macrophytes on the littoral ecosystem structure and function of a young Colorado Reservoir. *Aquatic Botany*, 85:37–43.
- Forum for Environment (2007). *Harnessing the Water Resources of Ethiopia for Sustainable Development in the New Ethiopian Millennium*, Addis Ababa, Ethiopia.
- Jacobus, V., Eshete, D., Abebe, G. and Leopold, A.J.N. (2011). The Composition of Fish Communities of nine Ethiopian Lakes along a North-South Gradient: Threats and Possible Solutions. *Animal Biology*.

- Klaudia, B. (2016). Accumulation of Cd and Pb in water, sediment and two littoral plants (*Phragmites australis*, *Typha angustifolia*) of freshwater ecosystem. *Archives of Environmental Protection. Institute of Environmental Engineering of the Polish Academy of Sciences, Zabrze, Poland.* 42(3): 47–57.
- Madsen, J.D. (2001). The interaction between water movement, sediment dynamics and submersed macrophytes. *Hydrobiologia*, 444: 71–84.
- Mulugeta, D. (2013). The Impact of Sedimentation and Climate Variability on the Hydrological Status of Lake Hawassa, South Ethiopia. PhD Thesis. Bonn University.
- Ottichilo, W.K. (2003). Use of Space Technology in Disaster Risk Assessment and Monitoring. 8 p.
- Padial, A.A., Bini, L.M. and Thomaz, S.M. (2008). The Study of Aquatic macrophytes in Neotropics: a Scientometrical View of the Main Trends and Gaps. *Brazilian Journal of Biology* 68(4, Suppl.): 1051–1059.
- Pattnaik, D.B.S.R. (2014). Species Diversity of Lake Hawassa, Ethiopia. *International Journal of Scientific Research*, 3(11): 35.
- Sagneeta, S. (2007). Role of Macrophytes in improving water quality of an aquatic ecosystem. *Journal of Applied Science and Environmental Management*, 11(4): 133–135.
- Small, C. and Cohen, J.E. (1999). Continental Physiography, Climate and the Global Distribution
- Tadesse, F. (2005). Trophic Analysis of Lake Hawassa using mass balance ecopath. Addis Ababa University School of Graduate Studies.
- Yezbie Kassa (2016). Macrophyte Ecology, Nutrient Dynamics and Water Quality of the Littoral Zone, and Yitamot Wetland, Lake Tana, Ethiopia. Department of Zoological Science. Addis Ababa University, Addis Ababa, Ethiopia.