

TREE HEIGHT – DIAMETER MODEL FOR THE INTERNATIONAL INSTITUTE OF AGRICULTURE (IITA) FOREST, IBADAN NIGERIA

***OLADOYE, A.O.,¹ AKINDELE, S.O.,² ADEKUNLE, V.A.J.² AND JOKODOLA, S.O.¹**

¹Department of Forestry and Wildlife Management, Federal University of Agriculture, Abeokuta, Nigeria

² Department of Forestry and Wood technology, Federal University of Technology, Akure, Nigeria

*Corresponding author: oladoyeao@funaab.edu.ng

Abstract

Tree height is one of the basic growth variables in forestry through which several other variables essential to forest management and decision-making and stand development over time are dependent. To this end, a study was conducted to develop models for estimating tree height- diameter relationship for IITA forest, Ibadan, Nigeria. Systematic sampling design was used to lay three straight line transects, four temporary plots of 0.25ha (50 m x 50 m) were laid in alternating position along each transect at 100 m interval to make up a total of 12 plots for the study, total height and Diameter at breast height (Dbh) were measured for all trees with Dbh \geq 10cm in every plot. A total of 299 trees, 46 species from 25 families were encountered. Mean Basal Area (Ba) ranged from 0.01 m² to 0.54m², mean volume ranged from 0.02 m³ to 11.93 m³, mean Dbh ranged from 10.00cm to 110 cm and mean total height ranged from 3.0 m to 34.40 m The Height- diameter relationship was modeled using R statistics. The data collected were split into two sets; the first set was the calibration set which was used for models' construction and the second was the validation set. and Akaike's Information Criterion (AIC), Bayesian Information Criterion (BIC) and Root Mean Square Error (RMSE) were the criteria used to select the best model. Based on these criteria, the power function ($H = 1.3 + 29.275BH^{0.724}$) was adjudged to be the most suitable model that best predict tree height for IITA forest because it has the lowest AIC and RMSE of 1387.598 and 5.616 respectively followed by Gompertz. Logistics and Ratkowsky has the same AIC and RMSE of 1394.859 and 5.697 respectively. T-test carried out to validate the best model at 95% probability level revealed that there was no significant difference ($P>0.05$) between the mean of predicted height (12.402m) and mean of the observed height (12.378m) with p-value of 0.976. However, close and low values of AIC and RMSE for all models tested suggest that all the models can adequately predict tree height for IITA forest and an indication that the models' estimators are efficient.

Keyword: Tree height, Diameter at Breast Height, Models, IITA, AIC

Introduction

Tree height is one of the most important variables crucial for quantitative assessment of forest stocks, however, it is difficult to measure, more time consuming, and expensive. In addition, the closed canopy nature of dense stands hinders visibility (Nguyen *et al.*, 2019, Sharma, Parton 2007; Colbert *et al.*, 2002; Staudhammer and Lemay, 2000) and others sources of errors and challenges to measuring tree height, especially in natural forest include visual obstructions, rounded crown forms, leaning trees and terrain (Mugasha *et al.*, 2013). Several other variables essential to forest management decision-making, such as tree scrolling and volume, stand dominant height and station quality index are derived from tree height, and the projection of stand development over time is based on precise height-diameter functions (Achille *et al.*, 2022; Baumeister, 2017; Calama and Montero, 2004).

Tree height measuring exercise is a tedious task and even more difficult in natural and complex tropical forests, where trees of various ages, species, sizes, vigor classes, crowns and shade tolerance levels coexist, than under uniform planting conditions and this could cause measurement errors (Calama and Montero, (2004); Fortin *et al.*, (2009); Zhang *et al.*, (2014), Temesgen *et al.* (2014) Maiti *et al.*, 2016) and Salami *et al.* (2021). Tree height-diameter models effectively describe the relationship between both tree characteristics at the stand level, and are frequently used in forest inventories for forest management and planning when sufficiently representative sample of diameter and height measurements are used (Ayoola and Silas, 2023). Information on tree

height and diameter is important for efficient management of trees in various ways such as to estimate the volume, the height of the trees in the forest which is also a pointer to evaluate site productivity and provides information on the competition of trees for nutrient, light and space within a stand (West, 2015, Ige *et al.*, 2019, Ayoola and Silas, 2023). Hence to save time and effort in forest inventory, it is important to develop tree height – diameter relationship models to predict height instead of direct measurement of height (Ahmadi *et al.*, 2013).

The International Institute of Tropical Agriculture (IITA) forest, Ibadan, Nigeria has been under protection since 1965 shortly after the whole area was acquired and has served as reservoir of numerous plant and animal species. A number of studies on regeneration dynamics Ariyo (2020), Salami and Akinyele (2018), Oladoye *et al.* (2014), Ariyo *et al.* (2013), socio-economics (Ariyo *et al.*, 2018.), yield models (Aghimien *et al.*, 2016), Wildlife (Osunsina *et al.*, 2012), there is no known or documented studies on tree height–diameter distribution model for IITA forest. Hence the objective of this study was to develop tree height–diameter relationship models for tree species in the IITA forest to promote easy estimation of tree heights.

Materials and Methods

Study Area

The study area is located on the one thousand hectares out of which 350 hectares IITA Forest Reserve has been under active protection since 1965 in the International Institute of Tropical Agriculture (I.I.T.A) campus at Idi-Ose, North of Ibadan, Nigeria on longitude 7° 30'N and latitude 3° 55' E and 243m above sea level. The site is within the humid

tropical lowland and has two distinct weather seasons: the longer wet season and shorter dry season. Wet season last for eight months and it extends from March to October while dry season last for four months from November to February. The rainfall pattern has bimodal peak with an annual total range between 1,300-1,500mm most of which falls between

May and September. The average daily temperature ranges between 21°C-23°C while the maximum is between 28°C and 34°C. Mean relative humidity is in the range of 64-84% (Ariyo, 2020, Aghiemien, *et al.*, 2016, Oladoye and Aduradola, 2015, Osunsina *et al.*, 2012, Tenkouano and Baiyeri, 2007).




Fig. 1: Map of the study Area

Study Design

Systematic sampling (line transect) technique was adopted for plot location in the reserve following Adekunle *et al* (2004), Oladoye *et al.*, (2020). A distance of 20 m from the forest boundary was measured to locate the first transect. The starting point of each transect was determined with the aid of Geographic Positioning System (GPS) receiver. Three transects were laid within the location and

four 0.25ha (50 m x 50 m) plots were laid in alternate position along each transect at 100 m interval to make a total of 12 plots for the study.

Data Collection

The data used for this study comprises tree height (H) and diameter at breast height (Dbh) for 299 trees from 46 species collected from twelve (12) temporary sample plots of 0.25 ha. Measurement and information collected are diameter at

breast height (Dbh) of diameter girth and tape Spiegel Relascope respectively. Also, trees were identified to species level.

Data Analysis

Tree height-diameter models that were used for this study as presented in table 1, are power, exponential, logistic, Gompertz, Ratkowsky and Modified Exponential, their consideration was based on their appropriate mathematical features such as (typical sigmoid shape, flexibility) and possible biological integration of parameters (eg. Upper asymptote, maximum or minimum growth rate as described by Huang (1999), Peng *et al.* (2001), Temesgen *et al.*, 2014; Mehtätalo *et al.*, 2015; Mensah *et al.*, 2018). The models selected were fitted by nonlinear least-squares (nls) estimations using the ‘nls’ function of the R Stats Package version 3.6.4 while the selection of the best fit model was based on the lowest RSE, mean bias, AIC and BIC values. Prior to model calibration, scattered plot of height against dbh of the data from the collected was plotted to know the relationship between the two variables across plots and appeared to be

nonlinear (Figure 2). Hence, the choice of nonlinear modelling. Residual analysis was conducted to identify the error structure. A weighting factor was found to be appropriate for achieving the equal error variance assumption to deal with heteroscedasticity of the data set.

Model validation is important before inferences deduced from them can be used with confidence. Validation involves the process of testing and comparing the models output with what is observed in the real world (Ige *et al.*, 2014).The data collected was randomly split into fitting (70%) and validation (30%) dataset following Akindele (1990), Ige *et al.*, (2019) Summary statistics of stand variables from sample plots used for model fitting and model validation are presented in table 2.

Fitting of Height-diameter Models

Prior to model calibration, scattered plot of total height against Dbh of the data from the field was plotted so as to know the relationship between the two variables and plots appeared to be nonlinear (Figures 2). Hence, the choice of using nonlinear modelling was considered.

Table 1: Tree Height–diameter models

| Non-Linear Height-Diameter Equations Selected | |
|--|--|
| Model | Equation |
| Exponential | $1.3 + ae^{bDbh}$1 |
| Modified Exponential | $1.3 + ae^{b/Dbh}$2 |
| Logistic | $H = 1.3 + a(1 + b(-cDBH))exp(Zeide,1993)$3 |
| Gompertz | $H = 1.3 + a \exp(\frac{b}{DBH} + c)$ (Ratkowsky,1990).....4 |
| Power | $H = 1.3 + aDBH^b$ (Arabatzis and Burkhart, 1992).....5 |
| Ratkowsky | $1.3 + \frac{a}{(1+e^{(b-c*Dbh)})}$6 |

Where H = Total Height (m), exp =Exponential, DBH = Diameter at Breast Height (cm), a, b, c = Parameter Estimates

Table 2: Descriptive statistics summary of the group data

| Variables | Fitting data (N trees = 220) | | | | Validation data (N trees = 79) | | | |
|----------------------|------------------------------|-------|------|--------|--------------------------------|-------|-------|--------|
| | Mean | SD | Min | Max | Mean | SD | Min | Max |
| DBH (cm) | 30.13 | 23.21 | 1.00 | 134.00 | 27.34 | 16.75 | 10.00 | 110.00 |
| THT (m) | 13.06 | 8.29 | 3.00 | 85.00 | 12.38 | 5.11 | 3.00 | 25.00 |
| VOL(m ³) | 1.57 | 4.83 | 0.02 | 50.46 | 0.84 | 1.58 | 0.02 | 10.97 |




Fig. 2: Relationship between Height and Diameter for IITA Forest, Nigeria

Assessment of Models

Models assessment was carried out to estimate recommendation of models with good fit for further uses: models’ examination at all stages of model design, fitting and implementation is important. T-test as described by Adekunle *et al.* (2013), Ige *et al.* (2019) was carried out to assessed the validity and suitability of the selected model by comparing the observed value for the dependent variable (height) with their predicted values. The decision rule for the test is to accept the null hypothesis (Ho: mean value observed = mean value predicted) if the p-value < 0.05.

The following statistical criteria were used.

Akaike’s Information Criterion (AIC)

This was computed using

$$AIC = n \times \ln\left(\frac{RSS}{n}\right) + 2 \times k \dots\dots 7$$

Bayesian Information Criterion (BIC)

$$BIC = \ln\left(\frac{RSS}{n}\right) + \frac{k}{n} \ln(n) \dots\dots 8$$

k = number of parameters in the model

n = number of regression coefficient

RSS being the residual sum of squares of the regression

Root Mean Square Error (RMSE)

$$RMSE = \sqrt{\sum \frac{(y_i - \hat{y}_i)^2}{n-p}} \dots\dots 9$$

Y_i =observed value of the dependent variable
 \hat{Y} =predicted value of the dependent variable
 n =number of observations
 p =number of parameters

Significance of Regression (F-ratio)

This was used to test the overall significance of the regression equation. The critical value of F (F-Tabulated) at equals 0.05 was compared with variance ratio F (F-calculated). When F-calculated is greater than the critical value (F-tabulated), such equation was considered significant and therefore adopted for prediction.

Results

A total, 299 trees were encountered representing 46 species from 25 families. *Elaeis guineensis* had the highest no of stem (30) followed by *Trilepisium madagascariense* (27), *Funtumia elastica*, *Holarrhena floribunda*, *Trichilia monadelphpha* had a total of 18 stems each. Mean Dbh ranged from 10.00cm to 110 cm and mean total height ranged from 3.0 m to 34.40 m Moraceae family had the highest number of species (6) while Sterculiaceae and Apocynaceae had four species each (table 3). Height-diameter relationship model for IITA forest was developed using six non-linear models and the parameter estimates and fit statistics presented in table 4. Total tree height and Diameter at breast height were used as dependent and independent variable respectively. Gompertz, logistics, power exponential, modified exponential and Ratkowsky were the models selected for developing the relationship. RMSE

AIC and BIC values ranged from 5.616 to 6.091, 1387.598 to 1423.346 and 1397.779 to 1433.527 respectively. Based on the AIC value and after evaluation and validation, power model produced the best fit for H-D for IITA forest. Power model also has the least RMSE (5.616) and BIC (1397.779). Modified exponential is adjudged to be the least model for developing H-D for IITA forest. Gompertz model is ranked second after power model with RMSE, AIC and BIC of 5.661, 1392.132, and 1405.706 respectively.

The result of T-test carried out to validate the best model at 95% probability level revealed that there was no significant difference ($P > 0.05$) between the mean of predicted height (12.402m) and mean of the observed height (12.378m) with p-value of 0.976. Figure 2 presented the scatter relationship between total height and Dbh at breast height very the actual field data with majority of the trees in the diameter range of 10-60cm. Figure 3 a-f presented the residual plot of the predicted height obtained from Gompertz, logistic, power, exponential, and modified exponential and Ratkowsky respectively. The figure indicated that a homogenous distribution of residuals and suggests that the assumption of regression was not violated by the models and is independent of experimental error. Figure 4 presented a graphically visualization of the fitted models comparison of all the six-nonlinear models which reveals that all the models have similar tendency to relatively estimate total height.

Table 3: Checklist of tree species, families and growth parameters of trees Encountered

| Species | | No of Stem | Mean DBH/cm | mean THT/m |
|------------------------------------|-----------------|------------|-------------|------------|
| <i>Aidia genipiflora</i> | Rubiaceae | 5 | 23.60 | 9.66 |
| <i>Albizia ferruginea</i> | Mimosaceae | 4 | 37.05 | 16.05 |
| <i>Albizia zygia</i> | Mimosaceae | 4 | 36.00 | 13.68 |
| <i>Alchornea laxiflora</i> | Euphorbiaceae | 1 | 12.00 | 5.50 |
| <i>Alstonia boonei</i> | Apocynaceae | 5 | 35.20 | 16.86 |
| <i>Antiaris africana</i> | Moraceae | 3 | 30.67 | 16.17 |
| <i>Antiaris toxicaria</i> | Moraceae | 6 | 55.00 | 13.47 |
| <i>Ceiba pentandra</i> | Bombacaceae | 10 | 65.90 | 25.49 |
| <i>Celtis philippensis</i> | Cannabaceae | 5 | 14.86 | 8.18 |
| <i>Celtis toka</i> | Ulmaceae | 6 | 15.67 | 7.42 |
| <i>Celtis zenkeri</i> | Ulmaceae | 10 | 23.00 | 13.92 |
| <i>Chrysophyllum albidum</i> | Sapotaceae | 7 | 17.14 | 9.36 |
| <i>Cleistopholis patens</i> | | 13 | 41.62 | 14.89 |
| <i>Cleistopholis philippensis</i> | Annonaceae | 2 | 26.00 | 12.00 |
| <i>Cola millenii</i> | Sterculiaceae | 5 | 16.20 | 9.58 |
| <i>Dialium guineensis</i> | Caesalpiniaceae | 1 | 12.00 | 11.00 |
| <i>Diospyros monbutensis</i> | Ebenaceae | 1 | 10.30 | 6.00 |
| <i>Elaeis guineensis</i> | Arecaceae | 30 | 36.00 | 13.95 |
| <i>Erythrina senegalensis</i> | Fabaceae | 3 | 19.67 | 12.67 |
| <i>Ficus exasperata</i> | Moraceae | 1 | 10.00 | 5.00 |
| <i>Ficus mucoso</i> | Moraceae | 1 | 110.00 | 20.00 |
| <i>Funtumia elastica</i> | Apocynaceae | 18 | 19.56 | 11.47 |
| <i>Holarrhena floribunda</i> | Apocynaceae | 19 | 30.44 | 15.28 |
| <i>Iecaniodiscus cupanioides</i> | Sapindaceae | 16 | 16.91 | 8.01 |
| <i>Lonchocarpus sericeus</i> | Papilionaceae | 2 | 26.67 | 11.27 |
| <i>Maesopsis emini</i> | Rhamnaceae | 7 | 43.14 | 20.31 |
| <i>Milicia excelsa</i> | Moraceae | 2 | 36.00 | 19.10 |
| <i>Millettia griffoniana</i> | Papilionaceae | 7 | 37.43 | 13.29 |
| <i>Myrianthus arboreus</i> | Moraceae | 1 | 28.00 | 7.00 |
| <i>Newbouldia laevis</i> | Boraginaceae | 3 | 22.75 | 5.25 |
| <i>Olex subscorpioidea</i> | Olacaceae | 1 | 10.00 | 6.00 |
| <i>Pouteria alnifolia</i> | Sapotaceae | 3 | 14.67 | 7.33 |
| <i>Ptercapous soyausii</i> | Fabaceae | 4 | 34.33 | 15.83 |
| <i>Pycnanthus angolensis</i> | Myristicaceae | 0 | 14.60 | 9.00 |
| <i>Ricinodendron heudelotii</i> | Euphorbiaceae | 1 | 54.00 | 19.00 |
| <i>Spathodea campanulata</i> | | 2 | 24.50 | 9.60 |
| <i>Spondias mombin</i> | Bignoniaceae | 15 | 21.25 | 11.18 |
| <i>Sterculia setigera</i> | Sterculiaceae | 8 | 23.38 | 14.46 |
| <i>Sterculia tragacantha</i> | Sterculiaceae | 9 | 16.51 | 9.81 |
| <i>Trichilia megalantha</i> | Meliaceae | 2 | 57.00 | 16.00 |
| <i>Trichilia monadelpha</i> | Meliaceae | 19 | 17.14 | 6.75 |
| <i>Trilepisium madagascariense</i> | | 27 | 36.36 | 13.76 |
| <i>Triplochiton scleroxylon</i> | Sterculiaceae | 3 | 65.33 | 34.40 |
| <i>Voacanga africana</i> | Apocynaceae | 1 | 18.00 | 3.70 |
| <i>Zanthoxylum leprieurii</i> | Rutaceae | 1 | 32.00 | 13.00 |
| <i>Zanthoxylum zantoloides</i> | Rutaceae | 5 | 19.67 | 14.95 |

Table 4: Regression Parameter estimates and fit statistics of Height-diameter relationship for IITA Forest

| S/N | Model Name | Model Parameter | | | Selection Criteria | | |
|-----|----------------------|-----------------|--------|-------|--------------------|----------|----------|
| | | a | b | c | RMSE | AIC | BIC |
| 1 | Gompertz | 43.906 | 2.289 | 1.822 | 5.661 | 1392.132 | 1405.706 |
| 2 | Logistic | 37.538 | 6.260 | 3.370 | 5.697 | 1394.859 | 1408.433 |
| 3 | Power | 29.275 | 0.724 | - | 5.616 | 1387.598 | 1397.779 |
| 4 | Exponential Modified | 7.552 | 1.315 | - | 5.955 | 1413.374 | 1423.555 |
| 5 | Exponential | 30.445 | -0.235 | - | 6.091 | 1423.346 | 1433.527 |
| 6 | Ratkowsky | 37.538 | 1.834 | 3.370 | 5.697 | 1394.859 | 1408.433 |




Fig. 3(a)




Fig. 3(b)




Fig. 3(c)




Fig. 3(d)




Fig. 3(e)




Fig. 3(f)

Figure 3: Residual plot using (a). Gompertz, (b). Logistic (c). Power, (d). Exponential, (e). Modified Exponential and (f). Ratkowsky for Height-Diameter Relationship in IITA Forest




Fig. 4: Graphical visualization of the fitted models comparison of all the six-nonlinear models

Discussion

Tree height and diameter at breast height, height of individual trees and stand are the two most commonly measures used variables in forest inventories for understanding the dynamics of forest

ecosystems describing forest structures, estimating stand, volume, biomass, Carbon and incorporating ecological characteristics of species constituting the canopy (Zhang *et al.*, 2014; Tsega *et al.*,

2018; Mensah *et al.*, 2018; Ng'andwe *et al.*, 2019; de Lima *et al.*, 2021)

Tree height diameter relationship for IITA forest was tested and evaluated with six nonlinear height diameter model functions in order to predict the height of trees using AIC, BIC and RMSE as the criteria to select the best models. Based on these criteria, the power function was adjudged to be the most suitable model that best predict tree height for IITA forest because it has the lowest AIC and RMSE of 1387.598 and 5.616 respectively followed by Gompertz. Logistics and Ratkowsky has the same AIC and RMSE of 1394.859 and 5.697 respectively. However, close and low values of AIC and RMSE for all models tested suggest that all the models can adequately predict tree height for IITA forest and an indication that the models estimators are efficient. This is similar to the findings of Ige *et al.* (2013) who reported low RMSE for Modified exponential, Oladoye *et al.*, (2015) Weibull model for *Nauclea diderrichii*, in Nigeria.

The power function has always been found suitable for predicting tree height in some tropical forest. Chenge, (2021) in Omo Strict nature reserve, Mensah *et al.* (2017) and (2018) in South African Misbelt forests and South Africa's indigenous forests respectively. Similar findings have been recorded by Imani *et al.*, (2017). The decision to use a model function to predict tree height using diameter at breast height as variable in natural forests may be a function of differences in tree height and diameter relationship in different forest sites, this is evidence in the findings of Banin *et al.*; (2012), Fayolle *et al.* (2016); Imani *et al.* (2017) and Chenge (2021).

The result of model validation showed that there was no significant difference

between mean of observed height (12.378) and mean of predicted height (12.402) with p – value of 0.97. This suggests the capability of the power model and other evaluated models in this study to effectively predict tree height and further validated the positive relationship between diameter at breast height and total tree height and the possibility of predicting tree height using diameter at breast height as predictor within the same location. (Moore *et al.*, 1996; Sharma, 2009; Avsar, 2004 and Osman *et al.*, 2013).

Residual plot for the six models selected showed a randomly distributed residuals with a zero mean and relatively homogenous variances except for exponential and modified exponential, respectively. The height diameter curve of power models indicates no violation of assumptions of non – linear regression similar to the finding of Chenge, (2021). The findings from this study further revealed the potentials of diameter at breast height to predict tree height based on the models statistics and the models' ability for predictive purposes. The results of these height curves in IITA forest revealed that the successful models will continue to perform very well within the range observed field data, prediction outside these range may produce different response by the same successful models.

Tree height diameter relationship can be influenced by a number of factors which can make the models to yield better convincing results and this include but no limited to stand density, climatic variations (Zheng *et al.* (2025), Salami *et al.* (2021); Kearsley *et al.* (2017), Shamaki, *et al.*, 2016) and this may explain the rationale behind the different model been adjudged for different forest reserves.

Conclusion

The relationship between height and diameter for trees in IITA forest was observed to follow a nonlinear pattern based on the preliminary scatter plots, hence the decision to carry out nonlinear modelling. Based on the different performance evaluation of the models, the power function, amongst all other tested functions, was considered most adequate for modeling height–diameter relationship of trees in IITA forest based on the AIC, BIC and RMSE value. Local height-diameter function which are only dependent on diameter at breast height can normally be applied to the location or locations with similar ecological conditions to where the data were collected. Hence the model developed and validated are adequate to provide more reliable estimate of height in the respective study area of area with similar ecological condition. Tree height diameter model developed in this study efficient and suitable for tree height estimation and could play a significant role in estimating tree biomass in IITA forest.

References

- Achille, A. N. Y., Beeckman, H., Din, N., Borgia, A. A. F., Christian, Z. J., and Marguerite, M. M. (2022). Height-Diameter Relationship of Some Forest Species Exploited for Wood in the Natural Tropical Forest of the Congo Basin. *Open Journal of Forestry*, 12: 235-247.
- Adekunle, V.A.J., Akindele, S.O. and Fuwape, J.A. (2004). Structure and yield models for tropical lowland rainforest ecosystem of South West Nigeria. *Food Agric Environ* 2: 395-399.
- Aghimien, E.V., Osho, J.S.A., Hauser, S., Deni, B. and Ade-Oni, V.D. (2016). Growth and Yield Models for Uneven-Aged Secondary Forest in IITA, Ibadan, Nigeria. *Forest Res.*, 5: 173. doi:10.4172/2168-9776.1000173
- Ahmadi, K., Alavi, S.J., Kouchaksaraei, M.T. and Aertsen, W. (2013). Non-linear height-diameter models for oriental beech (*Fagus orientalis* Lipsky) in the Hyrcanian forests, Iran. *Biotechnologie, Agronomie, Société et Environnement*, 17: 431–440.
- Akindele, S.O. (1990). Site quality assessment and yield equation for Teak plantation in the dry forest zone of Nigeria. Ph.D Thesis, Department of Forest Resources Management, University of Ibadan, Nigeria. 180p
- Arabatzis, A.A. and Burkhart, H.E. (1992). An evaluation of sampling methods and model forms for estimating height-diameter relationships in loblolly pine plantations. *Forest Science*, 38: 192–198.
- Ariyo, O.C. (2020). Comparative Analyses of Diversity and Similarity Indices of West Bank Forest and Block A Forest of the International Institute of Tropical Agriculture (IITA) Ibadan, Oyo State, Nigeria
- Ariyo, O.C., Okojie, L.O. and Ariyo, M.O. (2018). Villagers Willingness to Pay for Forest Conservation in Ibadan, Oyo State, Nigeria. *Asian Journal of Agricultural Extension, Economics and Sociology*, 23(4): 1–14.
- Avsar, M.D. (2004). The relationship between Diameter at breast height, tree height and crown diameter in Calabrain Pinus (*Pinus*

- brutia* Ten) of Baskonus Mountain, Kahramanmaras, Turkey. *Journal of Biological Science*, 4(4): 437-440.
- Ayoola J.O. and Silas A.E. (2023). Height Diameter Model for Selected Trees Within University of Ibadan, Oyo State, Nigeria. *Journal of Research in Forestry, Wildlife and Environment*, 15(1): 1-8 <http://www.ajol.info/index.php/jrfe>
- Banin, L., Feldpausch, T.R. and Phillips, O.L. (2012). What controls tropical forest architecture? Testing environmental, structural and floristic drivers. *Glob. Ecol. Biogeogr.* 21: 1179–1190. doi:10.1111/j.1466-8238.2012.00778.
- Baumeister, M. (2017). La dendrométrie ou les mathématiques du forestier. Arbres et forêts, Zimmer, L'Expertise Forestière.
- Calama, R. and Montero, G. (2004). Interregional Nonlinear Height-Diameter Model with Random Coefficients for Stone Pine in Spain. *Canadian Journal of Forest Research*, 34: 150-163.
- Chenge, I.B. (2021). Height–diameter relationship of trees in Omo strict nature forest reserve, Nigeria. *Trees, Forests and People*. (3). www.elsevier.com/locate/tfp
- Colbert, K.C., Larsen, D.R. and Lootens, J.R. (2002). Height–diameter equations for thirteen midwestern bottomland hardwood species. *Northern Journal of Applied Forestry*, 19: 171–176.
- de Lima, R.B., Rutishauser, E., da Silva, J.A.A., Guedes, M.C., Herault, B. and de Oliveira, C.P. (2021). Accurate Estimation of Commercial Volume in Tropical Forests *Forest Science*, 67: 14–21. <https://doi.org/10.1093/forsci/fxaa032>.
- Fayolle, A., Loubota Panzou, G.J., Drouet, T., Swaine, M.D., Bauwens, S., Vleminckx, J., Biwole, A., Lejeune, P. and Doucet, J.-L. (2016). Taller trees, denser stands and greater biomass in semi deciduous than in evergreen lowland central African forests. *Forest Ecology and Management*, 374: 42–50. <https://doi.org/10.1016/j.foreco.2016.04.033>
- Fortin, M., Bernier, S., Saucier, J.-P., and Labbé, F. 2009. Une relation hauteur-diamètre tenant compte de l'influence de la station et du climat pour 20 espèces commerciales du Québec. In Mémoire de recherche forestière n°153. Direction de la recherche forestière (41 p). Ministère des Ressources naturelles et de la Faune, Gouvernement du Québec.
- Huang, S. (1999). Ecoregion-based individual tree height-diameter models for lodgepole pine in Alberta. *Western Journal of Applied Forestry*, 14(4): 186–193.
- Ige, P.O., Adesoye, P.O. and Akinyemi, O.D. (2014). Modelling diameter distributions of a tropical natural forest in south west Nigeria with the beta distribution function. *Elixir Agriculture*, 74: 27160-27166.
- Ige, P.O., Akinyemi, G.O. and Smith, A.S. (2013). Nonlinear growth functions for modeling tree height–diameter relationships for *Gmelina arborea* (Roxb.) in south-

- west Nigeria, *Forest Science and Technology*, 9(1): 20-24.
- Ige, P.O. Adeoti, O.O and. Animashaun, Z.T. (2019). Non-Linear Height–Diameter Relationship Models for Four Accessions of *Parkia biglobosa* (Jacq. Benth) Plantation in Wasangare, Oyo State, Nigeria. *Journal of Agriculture and Environment*, 15(2): 181-194.
- Imani, G., Boyemba, F. and Lewis, S. (2017). Height-diameter allometry and above ground biomass in tropical montane forests: insights from the Albertine Rift in Africa. *PLoS One* 12: e0179653. doi:10.1371/journal.pone.0179653.
- Maiti, R., Rodriguez, H. and Kumari, A. (2016). Qualitative and Quantitative Characterization of Wood Fibers of Shrubs and Tree Species of the Tamaulipan Thorn Scrub, Northeastern Mexico and Its Possible Relation to Wood Quality and Utilization. *American Journal of Plant Sciences*, 7: 1046-1057.
- Mehtätalo, L., de-Miguel, S., and Gregoire, T. G. (2015). Modeling height–diameter curves for prediction. *Canadian Journal of Forest Research*, 45(7): 826–837.
- Mensah, S., Pienaar, O.L., Kunneke, A., du Toit, B., Seydack, A. and Uhl, E. (2018). Height—Diameter allometry in South Africa’s indigenous high forests: Assessing generic models performance and function forms. *Forest Ecology and Management*. 410: 1–11. <https://doi.org/10.1016/j.foreco.2017.12.030>.
- Moore, J.A., Zhang, L. and Stuck, D. 1996. Height-Diameter Equations for Ten Tree Species in the Inland Northwest. *Western Journal of Applied Forestry*, 11(4): 132–137. <https://doi.org/10.1093/wjaf/11.4.132>.
- Mugasha, W.A., Bollandås, O.M. and Eid, T. (2013). Relationships between diameter and height of trees in natural tropical forest in Tanzania. *Southern Forests*, 75: 221–237.
- Ng'andwe, P., Chungu, D., Yambayamba, A.M. and Chilambwe, A. (2019). Modeling the height-diameter relationship of planted *Pinus kesiya* in Zambia. *Forest Ecology and Management*, 447: 1-11.
- Nguyen Thanh, T., Tai Tien, D. and Shen, H.L. (2019). Height-diameter relationship for *Pinus koraiensis* in Mengjiagang Forest Farm of Northeast China using nonlinear regressions and artificial neural network models. *J. For. Sci.*, 65: 134–143.
- Oladoye, A.O. and Aduradola, A.M. (2015). Assessment of Floristic composition of Forest Undergrowth of International Institute of Tropical Agriculture (IITA) Forest Reserve Ibadan, Nigeria. *International Journal of Ecology and Environmental Studies*, 3(2): 74-84.
- Oladoye, A.O., Sodiya O. E., Ezenwenyi J. U., Basiru A.O, and Soaga J.A. 2015. Tree height prediction models for *Nauclea diderrichii* (De Wild. and Th.Dur.) Merrill Plantation in Area J4, Ogun State, Nigeria. *African Journal of Agriculture, Technology and Environment*, 4(2):46 – 60.
- Osman, E.I., Mamoun, H., Idris, El Zein, A., Ibrahim, E.I. and Mugira, M.

- (2013). Height-Diameter Prediction Models for Some Utilitarian Natural Tree species. *Journal of Forest Products and Industries*, 2013, 2(2): 31-39 ISSN 2325-453
- Osunsina, I.O.O., Inah, E. I., Ogunjinmi, A.A., Onadeko, S.A. and Osunsina, O. (2012). Distribution and Diversity of Flora and Fauna in International Institute of Tropical Agriculture (IITA) Forest and Nature Reserve, Ibadan. Oyo State. Nigeria. *Journal of Agriculture, Forestry and the Social Sciences* 10 (.2): 289-303.
- Peng, C., Zhang, L. and Liu, J. (2001). Developing and validating nonlinear height-diameter models for major tree species of Ontario's boreal forests. *Northern Journal of Applied Forestry*, 18: 87-94.
- Ratkowsky, D.A. (1990). Handbook of Nonlinear Regression, Marcel Dekker, New York, NY, USA.
- Salami, K.D. and Akinyele, A.O. (2018). Floristic Composition, Structure and Diversity Distribution in Omo Biosphere Reserve, Ogun State, Nigeria. *Ife Journal of Science*, 20(3): 639-648.
- Salami, K.D., Agbo-Adediran, A.O. and Kankomi, D.A. (2021). Development of Tree Height-Diameter Models for Arakanga Forest Reserve, Abeokuta, Ogun State. *Journal of Research in Forestry, Wildlife and Environment*, 13(1): 163-171.
- Shamaki, S.B., Akindele, S.O., Isah, A.D. and Mohammed, I. (2016) Height-Diameter relationship for Teak (*Tectona grandis*) Plantation in Nimbia Forest Reserve, Nigeria. *Asian Journal of Environment and Ecology*, 1(1): 1-7. DOI:10.9734/AJEE/2016/30635.
- Sharma M., and Parton J. 2007. Height-diameter equations for boreal tree species in Ontario using a mixed-effects modeling approach. *Forest Ecology and Management*, 249: 187–198.
- Sharma R.P. (2009). Modelling height-diameter relationships for Chir pine trees, *Banko Janakari*, 19:3–9.
- Staudhammer, C. and LeMay, V. (2000). Height prediction equations using diameter and stand density measures. *The Forestry Chronicle*, 76: 303–309.
- Temesgen, H. and Gadow, K.V. (2004). Generalized height–diameter models – an application for major tree species in complex stands of interior British Columbia. *European Journal of Forest Research*, 123: 45–51.
- Temesgen, H., Affleck, D., Poudel, K., Gray, A. and Sessions, J. (2014). A review of the challenges and opportunities in estimating tree height using height–diameter models. *Western Journal of Applied Forestry*, 29(1): 15–25.
- Tenkouano, K. and Baiyeri, P. (2007). Adoption pattern and yield stability of banana and plantain genotypes grown in contrasting agro- ecology zone in Nigeria,” *African Crop Science Conference Proceedings*, 8: 377–384.
- Tsega, M. Guadie, A., Teffera, Z.L., Belayneh, Y. and Niu, D. (2018). Development and validation of height-diameter models for *Cupressus lusitanica* in Gerged Forest, Ethiopia Forest

- Science and Technology*, 14(3): 138-144.
- West, P.W. (2015). Tree and forest measurement (3rd edition). Springer Cham, Heidelberg, New York.
- Zeide, B. (1993). Analysis of growth equation,” *Forest Science*, 39: 594–616, 1993.
- Zhang, X., Duan, A., Zhang, J. and Xiang, C. (2014). Estimating Tree Height-Diameter Models with the Bayesian Method. *Scientific World Journal* Volume 2014, Article ID 683691, 9 pages <http://dx.doi.org/10.1155/2014/683691>.
- Hindawi Publishing Corporation
- Zhang, L. (1997). Cross-validation of non-linear growth functions for modelling tree height-diameter relationships, *Annals of Botany*, 79(3): 251–257.
- Zheng, Y., Yue, Y., Gao, R. and Sharma, R. P. (2025). Developing mixed-effects height–diameter models using stand and climatic variables. *Scientific Reports*. The study demonstrated that stand density, mean annual precipitation, soil characteristics, and competition indices significantly influence tree height–diameter relationships.